

# CONCERTA<sup>®</sup> Extended-Release Tablets

## PRODUCT INFORMATION


**DRUG DEPENDENCE:** CONCERTA should be given cautiously to patients with a history of drug dependence or alcoholism. Chronic abusive use can lead to marked tolerance and psychological dependence with varying degrees of abnormal behaviour. Frank psychotic episodes can occur, especially with parenteral abuse. Careful supervision is required during withdrawal from abusive use since severe depression may occur. Withdrawal following chronic therapeutic use may unmask symptoms of the underlying disorder that may require follow-up.

### NAME OF THE MEDICINE

Methylphenidate hydrochloride

### DESCRIPTION

Methylphenidate hydrochloride is the racemic mixture of d,l-*threo*-methyl  $\alpha$ -phenyl-2-piperidineacetate hydrochloride. The d-(R,R)-isomer is pharmacologically more active than the l-(S,S)-isomer. Methylphenidate hydrochloride is a white, odourless crystalline powder. Its solutions are acid to litmus. It is freely soluble in water and in methanol, soluble in alcohol, and slightly soluble in chloroform and in acetone. It has the following structural formula:



C<sub>14</sub>H<sub>19</sub>NO<sub>2</sub>·HCl

MW 269.77

CAS-298-59-9 (methylphenidate hydrochloride)

CONCERTA is available as an extended-release tablet for once-a-day oral administration containing 18, 27, 36 or 54 mg methylphenidate hydrochloride. It is designed to have a 12-hour duration of effect.

CONCERTA tablets contain the following inactive ingredients: butylated hydroxytoluene, carnauba wax, cellulose acetate, hypromellose, lactose, Opacode black NS-78-17715, Opadry clear YS-1-19025-A, phosphoric acid, poloxamer, polyethylene oxide, povidone, sodium chloride, stearic acid, succinic acid and synthetic iron oxides. The 18 mg tablet also contains Opadry II yellow YS-30-12788-A. The 27 mg tablet also contains Opadry II grey Y-30-17528. The 36 mg tablet also contains Opadry II white Y-30-18037. The 54 mg tablet also contains Opadry II red Y-30-15567-A.

### PHARMACOLOGY

#### Pharmacodynamics

Methylphenidate is a central nervous system stimulant. The mode of therapeutic action in Attention Deficit Hyperactivity Disorder (ADHD) is not known. Methylphenidate is thought to block the reuptake of noradrenaline and dopamine into the presynaptic neuron and increase the release of these monoamines into the extraneuronal space.

#### Pharmacokinetics

**Absorption:** Methylphenidate is readily absorbed. Following oral administration of CONCERTA to adults, the drug overcoat dissolves and plasma methylphenidate concentrations increase rapidly reaching an initial maximum at about 1 to 2 hours. The methylphenidate contained in two internal drug layers is gradually released over the next few hours. Peak plasma concentrations are achieved at about 6 to 8 hours after which plasma levels of methylphenidate gradually decrease. CONCERTA once daily minimises the fluctuations between peak and trough concentrations associated with immediate-release methylphenidate three times daily. The extent of absorption of CONCERTA once daily is generally comparable to conventional immediate release preparations given three times daily.

Following the administration of CONCERTA 18 mg once daily in 36 adults, the mean pharmacokinetic parameters were  $C_{max}$   $3.7 \pm 1.0$  ng/mL,  $T_{max}$   $6.8 \pm 1.8$  h,  $AUC_{\infty}$   $41.8 \pm 13.9$  ngh/mL and  $t_{1/2}$   $3.5 \pm 0.4$  h. No differences in the pharmacokinetics of CONCERTA were noted following single and repeated once daily dosing indicating no significant drug accumulation. The AUC and  $t_{1/2}$  following repeated once daily dosing are similar to those following the first dose of CONCERTA 18 mg.

Following administration of CONCERTA in single doses of 18, 36 and 54 mg/day to adults,  $C_{max}$  and  $AUC_{inf}$  of d-methylphenidate were proportional to dose, whereas l-methylphenidate  $C_{max}$  and  $AUC_{inf}$  increased disproportionately with respect to dose. Following administration of CONCERTA, plasma concentrations of the l-isomer were approximately 1/40th the plasma concentrations of the d-isomer.

In healthy adults, single and multiple dosing of once daily CONCERTA doses from 54 to 144 mg/day resulted in linear and dose proportional increases in  $C_{max}$  and  $AUC_{inf}$  for total methylphenidate (MPH) and its major metabolite, (alpha)-phenyl-piperidine acetic acid (PPAA). The single dose and steady state (Day 4) clearance and half-life parameters were similar, indicating that there was no time dependency in the pharmacokinetics of methylphenidate. The ratio of metabolite (PPAA) to parent drug (MPH) was constant across doses from 54 to 144 mg/day, both after single dose and upon multiple dosing.

Pharmacokinetic equivalence has been demonstrated for two 27-mg CONCERTA tablets with three 18-mg CONCERTA tablets. The mean values of the treatment ratio (2 x 27 mg fasted/3 x 18 mg fasted) of the log-transformed pharmacokinetic values for  $C_{max}$ ,  $T_{max}$  and  $AUC_{inf}$  were 101.1%, 104.3% and 100.3% respectively. The 90% CIs for the treatment ratios were within the pre-specified 80% - 125% range.

Studies on the effects of dosing after overnight fasting, after consumption of a normal breakfast and a high-fat breakfast showed no differences in pharmacokinetics or pharmacodynamics of CONCERTA. There is no evidence of dose dumping in the presence or absence of food.

**Distribution:** Plasma methylphenidate concentrations in adults decline biexponentially following oral administration. The terminal plasma half-life of methylphenidate in adults following oral administration of CONCERTA was approximately 3.5 hours.

**Metabolism:** In humans, methylphenidate is metabolised primarily by de-esterification to  $\alpha$ -phenyl-piperidine acetic acid (PPAA) which has little or no pharmacologic activity. In adults the metabolism of CONCERTA once daily, as evaluated by metabolism to PPAA, is similar to that of methylphenidate three times daily. The metabolism of single and repeated once daily doses of CONCERTA is similar.

**Elimination:** After oral dosing of radiolabelled methylphenidate in humans, about 90% of the radioactivity was recovered in urine. The main urinary metabolite was PPAA, accounting for approximately 80% of the dose. Since renal clearance is not an important route of methylphenidate clearance, renal insufficiency is not expected to have a significant effect on the pharmacokinetics of CONCERTA.

## CLINICAL TRIALS

### Children

CONCERTA was demonstrated to be effective in the treatment of ADHD, in children aged 6 to 12 years, in three pivotal studies. Studies 1 and 2 were single-centre, double-blind, double-dummy, randomised, placebo and active-controlled, crossover comparisons (n = 64 and 70). Study 3 was a multicentre, 4 week, double-blind, double-dummy, randomised, placebo and active-controlled, parallel study (n = 282). The primary comparison of interest in all three trials was CONCERTA versus placebo.

The primary efficacy parameter for CONCERTA was the Inattention/Overactivity with Aggression (IOWA) Conners I/O subscale rated by the community school teacher. Statistically significant (p <

0.001) reduction in the Inattention/Overactivity subscale versus placebo was shown consistently across all three controlled studies for CONCERTA once daily.

Onset and duration of efficacy were assessed by the laboratory school teacher using the SKAMP (Swanson, Kotkin, Agler, M-Fynn and Pelham) combined attention ratings for studies 1 and 2. The onset of efficacy was estimated to be 1.5 hours and duration continued through to 12 hours. Patients demonstrated higher productivity and greater accuracy during CONCERTA treatment.

### **Adults**

Two double-blind, placebo-controlled studies were conducted in 627 adults aged 18 to 65 years. The controlled studies compared CONCERTA administered once daily and placebo in a multi-centre, parallel group, 5-week, fixed-dose study (Study 4) (18, 36, and 72 mg/day) and in a multi-centre, parallel group, 7-week dose-titration study (Study 5) (36 to 108 mg/day).

Study 4 was a multi-centre, double-blind, randomized, placebo-controlled, parallel group, dose-response study (5-week duration) with 3 fixed dose groups (18, 36, and 72 mg). Patients were randomized to receive CONCERTA administered at doses of 18 mg (n=101), 36 mg (n=102), 72 mg/day (n=102), or placebo (n=96). All three doses of CONCERTA were statistically significantly more effective than placebo in improving CAARS (Conners' Adult ADHD Rating Scale) total scores at double-blind end point in adult subjects with ADHD.

Study 5 demonstrated the effectiveness of CONCERTA in the treatment of ADHD in adults aged 18 to 65 years at doses from 36 mg/day to 108 mg/day based on the change from baseline to final study visit on the Adult ADHD Investigator Rating Scale (AISRS). Of 226 patients who entered the 7-week trial, 110 were randomized to CONCERTA and 116 were randomized to placebo. Treatment was initiated at 36 mg/day and patients continued with incremental increases of 18 mg/day (36 to 108 mg/day) based on meeting specific improvement criteria with acceptable tolerability. At the final study visit, mean change scores (LS Mean, SEM) for the investigator rating on the AISRS demonstrated that CONCERTA was statistically significantly superior to placebo.

## **INDICATIONS**

CONCERTA is indicated for the treatment of Attention Deficit Hyperactivity Disorder (ADHD). Treatment should be commenced by a specialist.

A diagnosis of Attention Deficit Hyperactivity Disorder (ADHD; DSM-IV) implies the presence of hyperactive-impulsive or inattentive symptoms that caused impairment and were present before age 7 years.

**Need for comprehensive treatment programme:** CONCERTA is indicated as an integral part of a total treatment program for ADHD that may include other measures (psychological, educational and social) for patients with this syndrome. Stimulants are not intended for use in the patient who exhibits symptoms secondary to environmental factors and/or other primary psychiatric disorders, including psychosis. Appropriate educational placement is essential and psychosocial intervention is often helpful. When remedial measures alone are insufficient, the decision to prescribe stimulant medication will depend upon the physician's assessment of the chronicity and severity of the patient's symptoms.

**Long term use:** The effectiveness of CONCERTA for long-term use, has not been systematically evaluated in controlled trials. Therefore the physician who elects to use CONCERTA for extended periods should periodically re-evaluate the long-term usefulness of the drug for the individual patient.

## **CONTRAINDICATIONS**

CONCERTA is contraindicated:

- in patients with known hypersensitivity to methylphenidate or any inactive ingredient used in this product (see **DESCRIPTION**);
- in patients with marked anxiety, tension, and agitation, since the drug may aggravate these symptoms;
- in patients with glaucoma;
- in patients with a family history or diagnosis of Tourette's syndrome;

- In combination with non-selective, irreversible monoamine oxidase (MAO) inhibitors / selective MAO-A inhibitors or within a minimum of 14 days following discontinuation of a non-selective, irreversible MAO inhibitor / selective MAO-A inhibitor (hypertensive crises may result);
- in patients with hyperthyroidism;
- in patients with severe angina pectoris;
- in patients with cardiac arrhythmia;
- in patients with phaeochromocytoma;
- In patients with known drug dependence or alcohol abuse;
- In patients with uncontrolled hypertension;
- In patients with ischaemic heart disease;
- In patients with myocardial infarctions;
- In patients who currently exhibit severe depression, anorexia nervosa, psychotic-symptoms or suicidal tendency, since CONCERTA might worsen these conditions.

## PRECAUTIONS

### Use with caution in the following circumstances

**Drug Dependence:** CONCERTA should be given cautiously to patients with a history of drug or alcohol dependence. Chronic abusive use can lead to marked tolerance and psychological dependence with varying degrees of abnormal behaviour. Careful supervision is required during withdrawal from abusive use since severe depression may occur. Withdrawal following chronic therapeutic use may unmask symptoms of the underlying disorder that may require follow-up

**Depression and Psychosis:** CONCERTA should not be used to treat severe depression or for the prevention or treatment of normal fatigue states. In psychotic patients administration of methylphenidate may exacerbate symptoms of behaviour disturbance and thought disorder.

Particular care should be taken in using stimulants to treat ADHD in patients with comorbid bipolar disorder because of concern for possible induction of a mixed/manic episode in such patients.

Treatment emergent psychotic or manic symptoms, e.g., hallucinations, delusional thinking or mania in patients without a prior history of psychotic illness or mania can be caused by stimulants at usual doses. If such symptoms occur, consideration should be given to a possible causal role of the stimulant, and discontinuation of treatment may be appropriate.

**Seizures:** There is some clinical evidence that methylphenidate may lower the convulsive threshold in patients with prior history of seizures, in patients with prior EEG abnormalities in absence of seizures, and very rarely in absence of history of seizures and no prior EEG evidence of seizures. In the presence of seizures, the drug should be discontinued.

**Potential for Gastrointestinal Obstruction:** CONCERTA tablet is non-deformable and does not appreciably change in shape in the GIT. It should not ordinarily be administered to patients with pre-existing severe GI narrowing (pathologic or iatrogenic) or in patients with dysphagia or significant difficulty in swallowing tablets. Due to the prolonged-release design of the tablet, CONCERTA should only be used in patients who are able to swallow the tablet whole.

**Visual Disturbance:** Difficulties with accommodation and blurring of vision have been reported with stimulant treatment.

**Motor and verbal tics:** CNS stimulants, including methylphenidate, have been associated with the onset or exacerbation of motor and verbal tics. Therefore, clinical evaluation for tics in patients should precede use of stimulant medication. Family history should be assessed.

### Sudden Death and Pre-existing Structural Cardiac Abnormalities or Other Serious Heart Problems

#### Children and Adolescents:

Sudden death has been reported in association with CNS stimulant treatment at usual doses in

children and adolescents with structural cardiac abnormalities or other serious heart problems. Although some serious heart problems alone carry an increased risk of sudden death, stimulant products generally should not be used in children or adolescents with known serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, or other serious cardiac problems that may place them at increased vulnerability to the sympathomimetic effects of a stimulant drug.

#### **Adults:**

Sudden deaths, stroke, and myocardial infarction have been reported in adults taking stimulant drugs at usual doses for ADHD. Although the role of stimulants in these adult cases is also unknown, adults have a greater likelihood than children of having serious structural cardiac abnormalities, cardiomyopathy, serious heart rhythm abnormalities, coronary artery disease, or other serious cardiac problems. Adults with such abnormalities should also generally not be treated with stimulant drugs.

It is essential that children, adolescents, or adults with pre-existing structural cardiac abnormalities or other serious heart problems being considered for treatment are assessed by a cardiologist before initiating treatment. Ongoing cardiologist supervision should be maintained throughout treatment in these patients.

### **Hypertension and Other Cardiovascular Conditions**

Children, adolescents or adults who are being considered for treatment with stimulant medications should have a careful history (including assessment for a family history of sudden death or ventricular arrhythmia) and physical exam to assess for the presence of cardiac disease, and should receive further cardiac evaluation if findings suggest such disease. Patients who develop symptoms such as exertional chest pain, unexplained syncope, or other symptoms suggestive of cardiac disease during stimulant treatment should undergo a prompt cardiac evaluation.

In the laboratory clinical trials in children, both CONCERTA and methylphenidate three times daily increased resting pulse by an average of 2 to 6 bpm and produced average increases of systolic and diastolic blood pressure of roughly 1 to 4 mm Hg during the day, relative to placebo. In placebo-controlled studies in adults, mean increases in resting pulse rate of approximately 4 to 6 bpm were observed with CONCERTA at endpoint vs. a mean change of roughly -2 to 3 bpm with placebo. Mean changes in blood pressure at endpoint ranged from about -1 to 1 mm Hg (systolic) and 0 to 1 mm Hg (diastolic) for CONCERTA and from -1 to 1 mm Hg (systolic) and -2 to 0 mm Hg (diastolic) for placebo. Therefore, caution is indicated in treating patients whose underlying medical conditions might be compromised by increases in blood pressure or heart rate. Blood pressure and cardiovascular status should be reviewed regularly during treatment with CONCERTA.

### **Aggression**

Aggressive behaviour or hostility is often observed in children and adolescents with ADHD, and has been reported in clinical trials and the postmarketing experience of some medications indicated for the treatment of ADHD. Although there is no systematic evidence that stimulants cause aggressive behaviour or hostility, patients beginning treatment for ADHD should be monitored for the appearance of or worsening of aggressive behaviour or hostility.

### **Haematologic Monitoring**

Periodic full blood count, differential and platelet counts are advised during prolonged therapy.

### **Renal Insufficiency**

There is no experience with the use of CONCERTA in patients with renal insufficiency. After oral administration of radiolabeled methylphenidate in humans, methylphenidate was extensively metabolized and approximately 80% of the radioactivity was excreted in the urine in the form of PPAA. Since renal clearance is not an important route of methylphenidate clearance, renal insufficiency is expected to have little effect on the pharmacokinetics of CONCERTA.

### **Hepatic Insufficiency**

There is no experience with the use of CONCERTA in patients with hepatic insufficiency.

### **Use in children**

The safety and efficacy of CONCERTA in children under 6 years old have not been established.

Long-term effects of methylphenidate in children are not yet available. Although a causal relationship has not been established, suppression of growth (ie weight gain and/or height) has been reported with the long-term use of stimulants in children. Therefore, patients requiring long-term therapy should be carefully monitored. Patients who are not growing or gaining weight as expected should have their treatment interrupted.

## **Carcinogenicity**

In a lifetime dietary carcinogenicity study carried out in mice, methylphenidate caused an increase in hepatocellular adenomas at a dose of 60–80 mg/kg/day, and in males only, an increase in hepatoblastomas (a relatively rare rodent malignant tumour type) at 60 mg/kg/day. These dose levels are approximately 3–8 fold the maximal recommended clinical dose on a mg/m<sup>2</sup> basis. There was no increase in tumours at 30–40 mg/kg/day (approximately 1-4 fold the maximal recommended clinical dose on a mg/m<sup>2</sup> basis). The mouse strain used is sensitive to the development of hepatic tumours, and the significance of these results to humans is not known. There was no evidence of carcinogenicity in two strains of transgenic mice administered methylphenidate in the diet for 24 weeks at doses up to 60–74 mg/kg/day (approximately 3–8 fold the maximal recommended clinical dose on a mg/m<sup>2</sup> basis) or in a lifetime dietary study in rats at doses up to 50 mg/kg/day (approximately 4–10 fold the maximal recommended clinical dose on a mg/m<sup>2</sup> basis).

## **Genotoxicity**

Methylphenidate was not mutagenic in the *in vitro* assays (Ames reverse mutation assay, mouse lymphoma cell forward mutation assay). Methylphenidate was weakly clastogenic *in vitro* (Chinese Hamster ovary cells) but was negative *in vivo* (mouse bone marrow micronucleus assay). Sister chromatid exchange assay results were positive only at high (cytotoxic) concentrations.

## **Effects on Fertility**

Dietary administration of methylphenidate to male and female mice at doses up to 150–160 mg/kg/day did not impair fertility in an 18-week continuous breeding study in which both parents and offspring were treated. This dose was approximately 7–16 fold the maximal recommended human dose on a mg/m<sup>2</sup> basis.

## **Use in pregnancy**

### **Category B3**

Oral administration of methylphenidate to rabbits during the period of organogenesis has produced teratogenic effects at doses of 200 mg/kg/day, associated with systemic exposure (plasma AUC) approximately 5 fold that in humans receiving the maximal recommended dose. The exposure at the no-effect dose in rabbits (60 mg/kg/day) was less than human exposure. Teratogenic effects were not seen in rats at oral methylphenidate doses up to 75 mg/kg/day, associated with systemic exposure of approximately 20 fold that in humans receiving the maximal dose. Oral administration of methylphenidate to rats from early pregnancy until weaning was associated with maternal toxicity, reduced offspring weight and marginal alterations in neuromotor performance in offspring at a maternal dose of 30 mg/kg/day, approximately 3-6 fold the maximum recommended clinical dose on a mg/m<sup>2</sup> basis. Oral administration of methylphenidate to juvenile male and female rats at doses of 12.5 mg/kg/day or greater from weaning through mating, pregnancy and lactation until offspring weaning was associated with reduced body weight gain and motor activity in males as well as reduced offspring weight and postnatal survival. The systemic exposure (plasma AUC) was 1 to 3 fold that expected in adults or children given the maximum recommended clinical dose, while the exposure at the no-effect dose was less than clinical exposure.

The safety of methylphenidate for use during human pregnancy has not been established, and no studies are available on the use of CONCERTA in pregnant women. CONCERTA should be used during pregnancy only if the potential benefit justifies the potential risk.

## **Use in lactation**

Oral administration of methylphenidate to rats from early pregnancy until weaning was associated with maternal toxicity, reduced offspring weight and marginal alterations in neuromotor performance in offspring at a maternal dose of 30 mg/kg/day, approximately 3-6 fold the maximum recommended clinical dose on a mg/m<sup>2</sup> basis. Oral administration of methylphenidate to juvenile male and female rats at doses of 12.5 mg/kg/day or greater from weaning through mating, pregnancy and lactation until

offspring weaning was associated with reduced body weight gain and motor activity in males as well as reduced offspring weight and postnatal survival. The systemic exposure (plasma AUC) was 1 to 3 fold that expected in adults or children given the maximum recommended clinical dose, while the exposure at the no-effect dose was less than clinical exposure.

Methylphenidate and / or its metabolites are excreted in milk in lactating rats at similar levels to plasma level. It is not known whether methylphenidate is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised if CONCERTA is administered to a nursing woman.

## **Interactions with other drugs**

Because of possible effects on blood pressure, CONCERTA should be used cautiously with pressor agents.

**Coumarin Anticoagulants, Antidepressants, and Selective Serotonin Reuptake Inhibitors:** Methylphenidate may inhibit the metabolism of coumarin anticoagulants, anticonvulsants (e.g. phenobarbital, phenytoin, primidone), and some antidepressants (tricyclics and selective serotonin reuptake inhibitors). Downward dose adjustment of these drugs may be required when given concomitantly with methylphenidate. It may be necessary to adjust the dosage and monitor plasma drug concentrations (or, in the case of coumarin, coagulation times), when initiating or discontinuing concomitant methylphenidate. Use of monoamine oxidase inhibitors is contraindicated, see CONTRAINDICATIONS

## **Effect on ability to drive or operate machinery**

**Stimulants may impair the ability of the patient to operate potentially hazardous machinery or vehicles. Patients should be cautioned accordingly until they are reasonably certain that CONCERTA does not adversely affect their ability to engage in such activities.**

## **Long-term Suppression of Growth**

Careful follow-up of weight and height in children aged 7 to 10 years who were randomized to either methylphenidate or non-medication treatment groups over 14 months, as well as in naturalistic subgroups of newly methylphenidate-treated and non-medication treated children over 36 months (to the ages of 10 to 13 years), suggests that consistently medicated children (i.e., treatment for 7 days per week throughout the year) have a temporary slowing in growth rate (on average, a total of about 2 cm less growth in height and 2.7 kg less growth in weight over 3 years), without evidence of growth rebound during this period of development.

Published data are inadequate to determine whether chronic use of amphetamines may cause similar suppression of growth, however, it is anticipated that they likely have this effect as well. Therefore, growth should be monitored during treatment with stimulants, and patients who are not growing or gaining height or weight as expected may need to have their treatment interrupted.

## **Instructions to the patient**

CONCERTA must be swallowed whole with the aid of liquids. Tablets should not be chewed, divided or crushed. The medication is contained within a nonabsorbable shell designed to release the drug at a controlled rate. The tablet shell, along with insoluble core components, is eliminated from the body; patients should not be concerned if they occasionally notice in their stool something that looks like a tablet.

## **ADVERSE EFFECTS**

### ***Clinical Trial Data***

### ***Double-Blind Data – Adverse Events Reported at ≥1% Frequency***

Adverse events (AEs) in either the paediatric or adult double-blind studies (Table 1 and 2) may be relevant for both patient populations.

### Paediatric Patients

The safety of CONCERTA was evaluated in 639 paediatric patients (children and adolescents) with ADHD who participated in 4 placebo-controlled, double-blind clinical trials. Three of the studies were conducted in children aged 6-12 years of age: two were cross-over studies in which patients received CONCERTA (doses of either 18 mg, 36 mg or 54 mg per day), immediate release methylphenidate and placebo for each of 7 days. The third study was a parallel group comparison in which patients were randomised to CONCERTA (doses of either 18 mg, 36 mg or 54 mg per day), immediate release methylphenidate or placebo for 28 days. In a fourth study, adolescents aged 13-18 years, receiving CONCERTA doses of 18 mg, 36 mg, 54 mg or 72 mg per day were randomised into a two week placebo-controlled, double-blind phase following an open-label 4 weeks titration phase. The information presented in this section was derived from pooled data.

Adverse Events (AEs) reported by  $\geq 1\%$  of CONCERTA-treated children and adolescents patients in these trials are shown in **Table 1**.

<b>Table 1. Adverse Events Reported by <math>\geq 1\%</math> of CONCERTA-Treated Children and Adolescents in 4 Placebo-Controlled, Double-Blind Clinical Trials</b>		
<b>System/Organ Class</b> Adverse Event	<b>CONCERTA</b> (n=321) %	<b>Placebo</b> (n=318) %
<b>Infections and Infestations</b>		
Nasopharyngitis	2.8	2.2
<b>Psychiatric Disorders</b>		
Insomnia*	2.8	0.3
<b>Nervous System Disorders</b>		
Headache	10.6	11.9
Dizziness	1.9	0
<b>Respiratory, Thoracic and Mediastinal Disorders</b>		
Cough	1.9	0.9
Oropharyngeal Pain	1.2	0.9
<b>Gastrointestinal Disorders</b>		
Abdominal Pain	6.2	3.8
Vomiting	2.8	1.6
<b>General Disorders</b>		
Pyrexia	2.2	0.9

\*Terms of Initial insomnia (CONCERTA=0.6%) and Insomnia (CONCERTA=2.2%) are combined into Insomnia

The majority of AEs were mild to moderate in severity.

### Adult Patients

The safety of CONCERTA was evaluated in 905 adult subjects with ADHD who participated in 3 placebo-controlled, double-blind clinical trials. The information presented in this section was derived from pooled data.

Adverse Events (AEs) reported by  $\geq 1\%$  of CONCERTA-treated adult subjects in these trials are shown in Table 2

<b>Table 2. Adverse Events Reported by <math>\geq 1\%</math> of CONCERTA-Treated Adults subjects in 3 Placebo-Controlled, Double Blind Trials</b>		
<b>System Organ Class</b> Adverse Event	<b>CONCERTA</b> N=596 (%)	<b>Placebo</b> N=309 (%)
<b>Cardiac disorders</b>		
Palpitations	4.5	0.6
Tachycardia	6.0	0.0

<b>Ear and labyrinth disorders</b>		
Vertigo	2.0	0.3
<b>Eye disorders</b>		
Vision blurred	1.3	1.0
Accommodation disorder	1.3	0.0
<b>Gastrointestinal disorders</b>		
Constipation	1.5	0.6
Dry mouth	15.1	3.6
Dyspepsia	2.0	1.9
Nausea	14.3	4.9
Vomiting	1.8	0.6
<b>General disorders and administration site conditions</b>		
Fatigue	4.7	4.2
Irritability	5.2	2.9
Thirst	1.8	0.6
Asthenia	1.2	0.0
<b>Infections and infestations</b>		
Influenza	8 (1.9)	4 (1.9)
Nasopharyngitis	19 (4.6)	11 (5.2)
Upper respiratory tract infection	1.7	1.0
Sinusitis	1.3	1.0
<b>Investigations</b>		
Blood pressure increased	2.5	1.9
Heart rate increased	3.0	1.9
Weight decreased	8.7	3.6
Alanine aminotransferase increased	1.0	0.0
<b>Metabolism and nutrition disorders</b>		
Anorexia	4.2	1.3
Decreased appetite	24.8	6.1
<b>Musculoskeletal and connective tissue disorders</b>		
Muscle tightness	1.3	0.0
Muscle spasms	1.0	0.3
Myalgia	5 (1.2)	3 (1.4)
<b>Nervous system disorders</b>		
Dizziness	7.4	5.5
Headache	24.2	18.8
Paraesthesia	5 (1.2)	0 (0.0)
Somnolence	7 (1.7)	9 (4.2)
Tension headache	1.0	0.3
Tremor	3.4	0.6
<b>Psychiatric disorders</b>		
Affect lability	1.3	0.6
Aggression	1.2	0.6

Agitation	3.2	0.6
Anxiety	8.4	2.9
Bruxism	1.5	0.6
Confusional state	1.0	0.3
Depressed mood	4.4	2.6
Depression	1.5	0.6
Initial insomnia	5.7	2.6
Insomnia	13.3	7.8
Libido decreased	1.3	0.6
Nervousness	2.3	0.6
Restlessness	4.0	0.0
Tension	1.3	0.3
Panic attack	1.3	0.3
<b>Respiratory, thoracic and mediastinal disorders</b>		
Cough	1.2	1.0
Oropharyngeal pain	1.5	1.3
Dyspnea	1.2	0.6
<b>Reproductive system and breast disorders</b>		
Erectile dysfunction	1.0	0.3
<b>Skin and subcutaneous tissue disorders</b>		
Hyperhidrosis	5.7	1.3
<b>Vascular disorders</b>		
Hypertension	2.2	1.6
Hot flush	1.3	0.6

The majority of AEs were mild to moderate in severity.

### ***Open-Label Data – Adverse Drug Reactions Reported at $\geq 1\%$ Frequency***

The safety of CONCERTA was evaluated in 3782 paediatric and adult patients with ADHD who participated in 12 open-label clinical trials. The information presented in this section was derived from pooled data.

Adverse Drug Reactions (ADRs) reported by  $\geq 1\%$  of CONCERTA-treated subjects in these trials and not listed in **Table 1 and 2** are shown in **Table 3**.

<b>Table 3. Adverse Drug Reactions Reported by <math>\geq 1\%</math> of CONCERTA-Treated Subjects in 12 Open-Label Clinical Trials</b>	
<b>System/Organ Class</b> Adverse Event	<b>CONCERTA</b> <b>(n=3782)</b> <b>%</b>
<b>General Disorders</b>	
Feeling jittery	1.4
<b>Psychiatric Disorders</b>	
Tic	2.0
Mood swings	1.1
<b>Skin and Subcutaneous Tissue Disorders</b>	
Rash	1.3
<b>Gastrointestinal disorders</b>	
Diarrhoea	2.4
Abdominal discomfort	1.3
Abdominal pain	1.2

The majority of ADRs were mild to moderate in severity.

***Double Blind and Open-Label Data – Adverse Drug Reactions Reported at <1% Frequency***

Additional ADRs that occurred in <1% of CONCERTA-treated paediatric and adult patients in the double-blind and open-label clinical datasets are listed in **Table 4**.

<b>Table 4. Adverse Drug Reactions Reported by &lt;1% of CONCERTA-Treated Paediatric and Adult subjects in Either Double-Blind or Open-Label Clinical Trials</b>
<b>Blood and Lymphatic System Disorders</b> Leukopenia
<b>Eye Disorders</b> Dry eye
<b>Investigations</b> Cardiac Murmur
<b>Nervous System Disorders</b> Lethargy, Somnolence, Psychomotor Hyperactivity, Sedation
<b>Psychiatric Disorders</b> Depression, Anger, Hypervigilance, Sleep Disorder, Mood Altered, , Agitation, Tearfulness,
<b>Skin and Subcutaneous Tissue Disorders</b> Rash, Rash-Macular

The majority of ADRs were mild to moderate in severity.

***Postmarketing Data***

Adverse events first identified as ADRs during postmarketing experience with CONCERTA are included in Table 5. The frequencies are provided according to the following convention:

Very common	≥1/10
Common	≥1/100 to <1/10
Uncommon	≥1/1,000 to <1/100
Rare	≥1/10,000 to <1/1,000
Very rare	<1/10,000, including isolated reports

<b>Table 5. Adverse Drug Reactions Identified During Postmarketing Experience with CONCERTA by Frequency Category Estimated from Spontaneous Reporting Rates</b>	
<b>Blood and Lymphatic System Disorders</b>	
<i>Very rare</i>	Pancytopenia, Thrombocytopenia, Thrombocytopenic purpura
<b>Psychiatric Disorders</b>	
<i>Rare</i>	Aggression
<i>Very rare</i>	Confusional state, Disorientation, Hallucination, Hallucination Auditory, Hallucination Visual, Mania, Nervousness, Restlessness
<b>Nervous System Disorders</b>	
<i>Very rare</i>	Convulsion, Grand Mal Convulsion, Dyskinesia
<b>Eye Disorders</b>	
<i>Very rare</i>	Diplopia, Mydriasis, Visual impairment
<b>Cardiac Disorders</b>	
<i>Very rare</i>	Angina Pectoris, Bradycardia, Extrasystoles, Supraventricular Tachycardia, Ventricular Extrasystoles
<b>Vascular Disorders</b>	
<i>Very rare</i>	Raynaud's Phenomenon
<b>Skin and Subcutaneous Tissue Disorders</b>	
<i>Very rare</i>	Alopecia, Erythema,
<b>Musculoskeletal, Connective Tissue and Bone Disorders</b>	
<i>Very rare</i>	Arthralgia, Myalgia, Muscle Twitching
<b>Immune System Disorders</b>	
<i>Rare</i>	Hypersensitivity reactions such as Angioedema, Anaphylactic reactions, Auricular swelling, Bullous conditions, Exfoliative conditions, Urticarias, Pruritus NEC, Rashes, Eruptions and Exanthemas NEC.
<b>General Disorders</b>	
<i>Rare</i>	Therapeutic Response Decreased
<i>Very rare</i>	Chest Pain, Chest Discomfort, Drug Effect Decreased, Hyperpyrexia
<b>Investigations</b>	
<i>Very rare</i>	Blood alkaline phosphatase Increased, Blood bilirubin Increased, Hepatic enzyme increased, Platelet count decreased, White blood cell count abnormal

Adverse events reported since market introduction in patients taking methylphenidate include suicide, suicide attempt, and suicide ideation. No causal relationship between methylphenidate and these events have been established.

## DOSAGE AND ADMINISTRATION

CONCERTA is administered orally once daily and should be taken in the morning.

CONCERTA must be swallowed whole with the aid of liquids, and must not be chewed, divided, or crushed.

CONCERTA may be administered with or without food.  
Treatment should be started on the lowest possible dose.

### **Children (greater than 6 years old) and adolescents:**

Dosage may be adjusted in 9 mg increments between 18 mg and 36mg and consecutively in an 18mg increment to a maximum of 54 mg/day taken once daily in the morning. In general, dosage adjustment may proceed at approximately weekly intervals.

### **Adults:**

Dosage can be adjusted from an initial dose of 18 or 36 mg/day in 18mg increments to a maximum of 72mg/day taken once daily in the morning. In general, dosage adjustment may proceed at approximately weekly intervals.

Patients respond at different dose levels and CONCERTA must be titrated to effect on an individual patient needs and response basis.

If treatment is restarted following discontinuation then dosing will need to be re-titrated rather than restarted from the previous dose. This approach should be considered for a discontinuation period of greater than 3 months.

Patients should be reviewed at least annually to assess if there is an ongoing requirement for treatment with CONCERTA. Blood pressure and cardiovascular status should also be regularly reviewed.

**Patients New to Methylphenidate:** The recommended starting dose of CONCERTA for patients who are not currently taking methylphenidate, or for patients who are on stimulants other than methylphenidate, is 18 mg once daily.

**Patients Currently Using Methylphenidate:** The recommended dose of CONCERTA for patients who are currently taking methylphenidate three times daily at doses of 15 – 60 mg per day is provided in **Table 6**.

**Table 6: Recommended dose conversions**

<b>Recommended CONCERTA dose</b>	<b>Previous methylphenidate dose</b>
18 mg once daily	5 mg methylphenidate three times daily
36 mg once daily	10 mg methylphenidate three times daily
54 mg once daily	15 mg methylphenidate three times daily
72 mg once daily	20 mg methylphenidate three times daily

Clinical judgement should be used when selecting the dose for patients currently taking methylphenidate in other regimens. If improvement is not observed after appropriate dosage adjustments over a one-month period, the drug should be discontinued.

### **Use in Infants and children**

Use of CONCERTA in patients under six years of age has not been studied in controlled trials. CONCERTA should not be used in patients under six years old.

### **Use in Elderly**

Use of CONCERTA in patients over 65 years of age has not been studied in controlled trials

### **OVERDOSAGE**

The prolonged release of methylphenidate from CONCERTA should be considered when treating patients with overdose.

### **Signs and Symptoms**

Signs and symptoms of CONCERTA overdose, resulting principally from overstimulation of the CNS and from excessive sympathomimetic effects, may include the following: vomiting, agitation, muscle

twitching, convulsions, grand mal convulsions, confusional state, hallucinations (auditory and/or visual), hyperhidrosis headache, pyrexia, tachycardia, palpitations, heart rate increased, sinus arrhythmias, hypertension, mydriasis, and dry mouth.

## **Treatment**

Treatment consists of appropriate supportive measures. The patient must be protected against self-injury and against external stimuli that would aggravate overstimulation already present. Measures to detoxify the gut include administration of activated charcoal and a cathartic. Intensive care must be provided to maintain adequate circulation and respiratory exchange; external cooling procedures may be required for hyperpyrexia.

Efficacy of peritoneal dialysis or extracorporeal haemodialysis for CONCERTA overdose has not been established.

## **PRESENTATION AND STORAGE**

CONCERTA 18 mg are yellow capsule-shaped tablets, with "alza 18" printed in black ink on one side.

CONCERTA 27 mg are grey capsule-shaped tablets, with "alza 27" printed in black ink on one side.

CONCERTA 36 mg are white capsule-shaped tablets, with "alza 36" printed in black ink on one side.

CONCERTA 54 mg are brownish-red capsule-shaped tablets, with "alza 54" printed in black ink on one side.

CONCERTA tablets are supplied in HDPE bottles containing silica gel desiccant.

Store below 25°C. Keep container tightly closed.

## **NAME AND ADDRESS OF THE SPONSOR**

JANSSEN-CILAG Pty Ltd  
1-5 Khartoum Road Macquarie Park NSW 2113 Australia  
NZ Office: Auckland New Zealand

## **POISON SCHEDULE OF THE MEDICINE**

S8 – Controlled Drug

**Date of TGA approval:** 4 November 2011